

User Meeting 2020

Contribution ID: 219

Type: Oral

Resonant Tender X-ray Diffraction for Disclosing the Molecular Packing of Paracrystalline Conjugated Polymer Films

Thursday, 19 November 2020 11:20 (20 minutes)

The performance of optoelectronic devices based on conjugated polymers is critically dependent upon molecular packing; however the paracrystalline nature of these materials limits the amount of information that can be extracted from conventional X-ray diffraction. Resonant diffraction (also known as anomalous diffraction) occurs when the X-ray energy used coincides with an X-ray absorption edge in one of the constituent elements in the sample. The rapid changes in diffraction intensity that occur as the X-ray energy is varied across an absorption edge provide additional information that is lost in a conventional nonresonant experiment. Taking advantage of the fact that many conjugated polymers contain sulfur as heteroatoms, this work reveals pronounced resonant diffraction effects at the sulfur K-edge with a particular focus on the well-studied electron transporting polymer poly([N,N -bis(2-octyldodecyl)-naphthalene-1,4,5,8-bis(dicarboximide)-2,6-diyl]-alt-5,5 -(2,2 -bithiophene)), P(NDI2ODT2). The observed behavior is found to be consistent with the theory of resonant diffraction, and by simulating the energy-dependent peak intensity based on proposed crystal structures for P(NDI2OD-T2) it is shown that resonant diffraction can discriminate between different crystalline packing structures. The utilization of resonant diffraction opens up a new way to unlock important microstructural information about conjugated polymers for which only a handful of diffraction peaks are typically available.

Primary author: Prof. MCNEILL, Chris (Monash University)

Presenter: Prof. MCNEILL, Chris (Monash University)

Session Classification: Session 2 - Advanced Materials & Hard Matter

Track Classification: Advanced Materials and Hard Matter