

Morphological transitions in aqueous surface active ionic liquid influenced by additives investigated through SANS

Friday, 13 November 2020 10:30 (20 minutes)

Designing microscopic interactions judiciously in the amphiphilic systems with added additives is advantageous in order to design various morphologies. Apart from the biological relevance, ease of formation, and long-term stability, cationic systems could form various morphologies including vesicles, which are the best alternatives to liposomes and niosomes and are potentially the best ever cavity to carry the drug molecules within its cage/bilayer. In a quest for the best cationic system to sequester functional molecules, herein we have investigated the concentration/composition induced morphological transitions in the cationic systems comprising of surface-active ionic liquids (SAILs) and various inorganic and organic electrolytes. The variables studied for the phase transition and the shape and size of the aggregates includes; alkyl chain length and head group structures of the SAILs and the concentration and structure of the additives. We used various spectroscopic, microscopic and scattering techniques to determine the size and shape of the aggregates including SANS. The micelle to vesicle transition through added additive is investigated by SANS. The size of micelle, vesicle and bilayer thickness of vesicle is also determined through SANS. For the first time, we used fluorescence resonance energy transfer (FRET) to determine the size of the aggregates and prove the significance of this technique by correlating data with SANS data. This work could be of interest in the field of drug delivery as the possible new nanocarriers.

Level of Expertise

Speakers Gender

Female

Do you wish to take part in the poster slam

Yes

Primary authors: Mrs JAIN, Monika; Mr MALEK, Naved

Presenter: Mrs JAIN, Monika

Session Classification: Advanced Materials

Track Classification: Advanced Materials