

Scaling behaviour of the skyrmion phases of Cu₂OSeO₃ single crystals from small angle neutron scattering

Thursday, 12 November 2020 17:40 (1 minute)

A skyrmion is a topological stable particle-like object comparable to a spin vortex at the nanometre scale. It consists of an about 50 nm large spin rotation and its spin winding number is quantised. Skyrmions emerge in chiral crystals as the result of competing symmetric exchange and asymmetric Dzyaloshinskii-Moriya (DM) interactions and typically form two dimensional hexagonal lattices perpendicular to an applied magnetic field. Its dynamics has links to flux line vortices as in high-temperature superconductors [1-2].

Cu₂OSeO₃ is a unique case of a multiferroic material where the skyrmion dynamics could be controlled through the application of an external electric field. The direct control of the skyrmion dynamics through a non-dissipative method would offer technological benefits applicable in energy-efficient data storage and data processing devices or for testing fundamental theories also related to the Higgs Boson whose theoretical description has similarities to skyrmions [3].

The technological applications crucially depend on the stability conditions of the skyrmion phase up to room temperature. While some materials host skyrmion lattices above room temperature [3], Cu₂OSeO₃ is the only insulating skyrmion material discovered so far, which orders magnetically below 58 K. It is interesting to note that the appearance of two different skyrmion phases have been reported along the temperature and magnetic field phase diagram of Cu₂OSeO₃ when the sample is aligned with its main crystallographic axes parallel to the incoming neutron beam and performing Zero Field Cooling (ZFC) or Field Cooling (FC) across the high-temperature skyrmion phase. However, the stabilisation processes of these two phases and their thermodynamic connection are still under debate [4-6]. We have used small angle neutron scattering and Lorentz transmission electron microscopy [7] to study the scaling behaviour of helical phase and the magnetic skyrmion lattices, i.e. the systematic change of their distances in single crystals of Cu₂OSeO₃ in order to gain insight on the balance between the different competing magnetic exchange interactions.

Therefore, we have examined the field, temperature and sample alignment dependence of the scaling behaviour of skyrmions as an order parameter for the emergence of the two aforementioned skyrmion phases. The obtained data provide valuable information on the formation mechanism of the skyrmions and their stability range. This is an important step towards the understanding of the manipulation of skyrmions, which is required for technological applications.

- [1] S. Muehlbauer et al., Science 323, 915 (2009)
- [2] S. Seki, X. Z. Yu, S. Iihwata, and Y. Tokura, Science 336, 198 (2009)
- [3] A. Fert, N. Reyren, V. Cros, Nat. Rev. Mats. 2, 01731 (2017)
- [4] A. Chacon, L. Heinen et al., Nat. Phys. 14, 936-941 (2018)
- [5] F. Qian, L. J. Bannenberg et al., Sci. Adv. 4, eaat7323 (2018)
- [6] L. J. Bannenberg, H. Wilhelm et al., npj Quantum Mater. 4, 11 (2019)
- [7] M.-G. Han, J. Garlow, Y. Kharkov, L. Camacho, R. Rov, G. Vats, J. Saucedo Flores, K. Kisslinger, T. Kato, O. Sushkov, Y. Zhu, C. Ulrich, T. Soehnel, and J. Seidel, Sci. Adv. 6, eaax2138 (2020)

Level of Expertise

Student

Speakers Gender

Male

Do you wish to take part in the poster slam

No

Primary author: Mr SAUCEDA FLORES, Jorge Arturo (School of Physics, University of New South Wales, Sydney 2052, Australia)

Co-authors: Ms ROV, Rosanna (School of Chemical Sciences, University of Auckland, Auckland 1142, New Zealand); Mr PERVEZ, Md. Firoz (School of Physics, UNSW Sydney, Sydney NSW 2052, Australia); SPASOVSKI, Martin (University of Auckland); O'BRIEN, Joel; VELLA, Joseph (The University of Auckland); Prof. SEIDEL, Jan (School of Material Science and Engineering, The University of New South Wales); YICK, Samuel (UNSW and ANSTO); Prof. GILBERT, Elliot (Bragg Institute, Australian Nuclear Science and Technology Organisation); Dr TRETIAKOV, Oleg (School of Physics, UNSW Sydney, Sydney NSW 2052, Australia); SOEHNEL, Tilo (The University of Auckland); ULRICH, Clemens (The University of New South Wales)

Presenter: Mr SAUCEDA FLORES, Jorge Arturo (School of Physics, University of New South Wales, Sydney 2052, Australia)

Session Classification: Poster Session

Track Classification: Magnetism & Condensed Matter